



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

or both  $a$  and  $b > \$1$ . Then as  $x$  is an integer, as  $a > \$1$ , and as B paid but \$6 for the first month's service,  $x \geq 5$ . Similarly, as  $y$  is an integer,  $b > \$1$ , and as C paid but \$7.20 for the second month's service  $(y+1) \geq 7$ , or  $y \geq 6$ . Hence as  $n=3+x+y$ ,  $n \geq 14$ . But as  $c=m/(n+6)=1$ ,  $m=(n+6)$ , whence  $m \geq \$20$ . But as not less than 5 horses were placed in the pasture the first month,  $m = \geq 11$ .

B had  $x$  horses in the pasture the first month, the rate then being  $m/n$ , or  $m/(m-6)$ , and paid \$6 for the service. Hence,  $xm/(m-6)=36$ , or reducing and transposing,  $(6-x)m=36$ , but as  $(6-x)=$ integer, and  $11 \leq m \leq 20$  (remembering that C had some horses in the pasture the first month, and that  $n=m-6$ ).  $6-x=2$ ,  $m=\$18$ , whence  $x=4$  and  $n=12$ . But  $y=n-3-x=5$ . As  $a=m/n=\$1.50$ ,  $b=m/(n+3)=\$1.20$ , and  $c=\$1$ , we readily find that

A owed \$4.50 for the first, and \$4.80 for the second month's service = \$9.30 total.

B owed \$6 for the second, and \$6 for the third month's service = \$12 total.

C owed \$7.50 for the first, and \$7 for the third month's service = \$14.50 total.

**II. Solution by REV. J. H. MEYER, S. J., Professor of Mathematics, College of the Sacred Heart, Augusta, Ga.; by M. R. BECK, Cleveland, Ohio, and by J. EDWARD SANDERS, Reinersville, Ohio.**

Using  $r$  as the number of dollars in one month's rent;  $x$  and  $y$  as the number of horses B and C put in at first, respectively, we obtain

$$\frac{rx}{3+x+y}=6, \text{ or } r=\frac{18+6x+6y}{x}. \quad (1)$$

$$\frac{r(y+1)}{6+x+y}=7.20, \text{ or } r=\frac{48.2+7.2x+7.2y}{y+1}. \quad (2)$$

$$\frac{5r}{9+x+y}=5, \text{ or } r=\frac{45+5x+5y}{5}. \quad (3)$$

Equating (1) and (3),  $90-15x+30y=5x^2+5xy \dots (4)$ . Equating (2) and (3),  $171+31x-14y=5xy+5y^2 \dots (5)$ . Adding (4) and (5),  $-5(x+y)^2+16(x+y)=-261 \dots (6)$ . Solving (6) for  $x+y$ ,  $x+y=9 \dots (7)$ . Substituting (7) in (3), (2), (1),  $r=18$ ,  $y=5$ ,  $x=4$ . For first month, \$1.50, for the second, \$1.20, and for the third, \$1.00 must be paid for each horse.

Hence, A owed \$9.30; B, \$12.00; C, \$14.50.

Also solved by O. L. Callecot, A. H. Holmes, L. E. Newcomb, J. Scheffer, G. B. M. Zerr, and the Proposer.

257. Proposed by L. E. NEWCOMB, Los Gatos, Cal.

Solve (1)  $x+y=10$ , (2)  $3x=\log_{10}y$ .

**I. Solution by HENRY HEATON, Belfield, N. D.**

We have  $3x=\log_{10}(10-x)$ , or  $10-x=10^{3x}$ . If we suppose  $x=\frac{1}{3}$ , we obtain  $9\frac{2}{3}=10$ . An error of  $\frac{1}{3}$ . If we suppose  $x=\frac{1}{4}$ , we get  $9.25=4.38$ . An error of 4.87. By position we obtain  $x=.328$ . Substituting this we get  $9.672=9.639$ . An error of -.043. Substituting .329 for  $x$  we get  $9.671=9.704$ . An error of

.033. By position the second time we have  $x=.3285$ . Substituting this, we obtain  $9.6715=9.6716$ . An error of .0001.

Hence  $x=.3285-$ , and  $y=10-.3285=9.6715+$ .

II. Solution by A. H. HOLMES, Brunswick, Maine, and by J. SCHEFFER, Hagerstown, Md.

$$x+y=10 \dots (1), \quad 3x=\log_{10}y \dots (2).$$

$$\therefore \log y=\log(10-x) \text{ and } \log_{10}y=M\log(10-x). \quad \therefore \log(10-x)=3x/M.$$

$$\log(10-x)=\frac{1}{M}+\log\left(1-\frac{x}{10}\right)=\frac{1}{M}-\frac{x}{10}-\frac{x^2}{200}-\frac{x^3}{3000}-\text{etc.}=\frac{3x}{M}$$

The series converges very rapidly and the first two terms are sufficient for a correct value of  $x$  to the fourth decimal figure.

$$\text{Hence } \frac{1}{.434294}-\frac{x}{10}-\frac{x^2}{200}=\frac{3x}{.434294}.$$

$$\therefore x=.3285-, \text{ and from (1), } y=9.6714+.$$

III. Solution by S. A. COREY, Hiteman, Iowa.

From (1),  $y=10-x$ , and substituting in (2),  $\log_{10}(10-x)=3x \dots (3)$ ,

$$\text{whence } \log_{10}[10(1-\frac{x}{10})]=3x \dots (4), \text{ and } \log_{10}10+\log_{10}(1-\frac{x}{10})=3x \dots (5).$$

$$\log_{10}(1-\frac{x}{10})=3x-1 \dots (6). \quad \text{But as } \log_{10}(1-\frac{x}{10})<0, \text{ we have } 3x<1. \quad \text{Let } 3x=1-3v. \quad \text{By substituting in (6), } \log_{10}(\frac{29}{30}+\frac{v}{10})=-3v, \text{ or } \log_{10}[\frac{29}{30}(1+\frac{3v}{29})]=-3v, \quad \log_{10}(1+\frac{3v}{29})=\log_{10}30-\log_{10}29-3v=a-3v \quad (a=.014,723,256,8) \dots (7).$$

$$\text{But } \log_{10}(1+\frac{3v}{29})>0, \text{ hence } 3v<a. \quad \text{Next let } 3v=a-3w. \quad \text{By substituting in (7), } \log_{10}(\frac{29+a}{29}-\frac{3w}{29})=3w, \quad \log_{10}\left[\left(\frac{29+a}{29}\right)\left(1-\frac{3w}{29+a}\right)\right]=3w. \quad \text{Whence } \log_{10}\left(1-\frac{3w}{29+a}\right)=3w+\log_{10}29-\log_{10}(29+a)=3w-.000,220,434,6 \dots (8).$$

$$\text{But } \log_{10}\left(1-\frac{3w}{29+a}\right)=\frac{-3mw}{29+a} \text{ nearly, where } m=\text{modulus of common system of logarithms, and therefore (8) becomes } \frac{3mw}{29+a}+3w=.000,220,434,6, \text{ nearly.}$$

$$\text{Whence, } w=.000,072,394,6, \text{ nearly; and } x=\frac{1}{3}-a/3+w=.328,497,975,7, \\ y=10-x=9.671,502,024,3.$$

Also solved by G. W. Greenwood, G. B. M. Zerr, and the Proposer.